< Share

PDF Tools

PDF

#### CA: A Cancer Journal for Clinicians / Volume 0, Issue 0

Original Article 🔂 Free Access

#### Cancer statistics, 2019

Rebecca L. Siegel MPH **▼**, Kimberly D. Miller MPH ... See all authors →

First published: 08 January 2019 https://doi.org/10.3322/caac.21551

About Sections

### Abstract

Each year, the American Cancer Society estimates the numbers of new cancer cases and deaths that will occur in the United States and compiles the most recent data on cancer incidence, mortality, and survival. Incidence data, available through 2015, were collected by the Surveillance, Epidemiology, and End Results Program; the National Program of Cancer Registries; and the North American Association of Central Cancer Registries. Mortality data, available through 2016, were collected by the National Center for Health Statistics. In 2019, 1,762,450 new cancer cases and 606,880 cancer deaths are projected to occur in the United States. Over the past decade of data, the cancer incidence rate (2006-2015) was stable in women and declined by approximately 2% per year in men, whereas the cancer death rate (2007-2016) declined annually by 1.4% and 1.8%, respectively. The overall cancer death rate dropped continuously from 1991 to 2016 by a total of 27%, translating into approximately 2,629,200 fewer cancer deaths than would have been expected if death rates had remained at their peak. Although the racial gap in cancer mortality is slowly narrowing, socioeconomic inequalities are widening, with the most notable gaps for the most preventable cancers. For example, compared with the most affluent counties, mortality rates in the poorest counties were 2-fold higher for cervical cancer and 40% higher for male lung and liver cancers during 2012-2016. Some states are home to both the wealthiest and the poorest counties, suggesting the opportunity for more equitable dissemination of effective cancer prevention, early

detection, and treatment strategies. A broader application of existing cancer control knowledge with an emphasis on disadvantaged groups would undoubtedly accelerate progress against cancer.

## Introduction

Cancer is a major public health problem worldwide and is the second leading cause of death in the United States. In this article, we provide the estimated numbers of new cancer cases and deaths in 2019 in the United States nationally and for each state, as well as a comprehensive overview of cancer occurrence based on the most current population-based data for cancer incidence through 2015 and for mortality through 2016. We also estimate the total number of deaths averted because of the continuous decline in cancer death rates since the early 1990s and analyze cancer mortality rates by county-level poverty.

# **Materials and Methods**

### Incidence and Mortality Data

Mortality data from 1930 to 2016 were provided by the National Center for Health Statistics (NCHS).1-3 Forty-seven states and the District of Columbia met data quality requirements for reporting to the national vital statistics system in 1930, and Texas, Alaska, and Hawaii began reporting in 1933, 1959, and 1960, respectively. The methods for abstraction and age adjustment of historic mortality data are described elsewhere.3, 4 Five-year mortality rates (2011-2015) for Puerto Rico were previously published in volume 3 of the North American Association of Central Cancer Registries' (NAACCR's) *Cancer in North America: 2011-2015*.5

Population-based cancer incidence data in the United States have been collected by the National Cancer Institute's (NCI's) Surveillance, Epidemiology, and End Results (SEER) Program since 1973 and by the Centers for Disease Control and Prevention's (CDC's) National Program of Cancer Registries (NPCR) since 1995. The SEER program is the only source for historic population-based incidence data. Long-term (1975–2015) incidence and survival trends were based on data from the 9 oldest SEER areas (Connecticut, Hawaii, Iowa, New Mexico, Utah, and the metropolitan areas of Atlanta, Detroit, San Francisco–Oakland, and Seattle–Puget Sound), representing approximately 9% of the US population.6, 7 The lifetime probability of developing cancer and contemporary stage distribution and survival statistics were based on data from all 18 SEER registries (the SEER 9 registries plus Alaska Natives, California, Georgia, Kentucky, Louisiana, and New Jersey), covering 28% of the US population.8 The probability of developing cancer was calculated using NCI's DevCan software (version 6.7.6).9 Some of the statistical information presented herein was adapted from data previously published in the *SEER Cancer Statistics Review 1975-2015*.10 The NAACCR compiles and reports incidence data from 1995 onward for registries that participate in the SEER program and/or the NPCR. These data approach 100% coverage of the US population for the most recent years and were the source for the projected new cancer cases in 2019 and cross-sectional incidence rates by state and race/ethnicity.11, 12 Some of the incidence data presented herein were previously published in volumes 1 and 2 of *Cancer in North America: 2011-2015*.13, 14

All cancer cases were classified according to the *International Classification of Diseases for Oncology* except childhood and adolescent cancers, which were classified according to the International Classification of Childhood Cancer (ICCC).15, 16 Causes of death were classified according to the *International Classification of Diseases*.17 All incidence and death rates were age standardized to the 2000 US standard population and expressed per 100,000 population, as calculated by NCI's SEER\*Stat software (version 8.3.5).18 The annual percent change in rates was quantified using NCI's Joinpoint Regression Program (version 4.6.0).19

Whenever possible, cancer incidence rates were adjusted for delays in reporting, which occur because of a lag in case capture or data corrections. Delay-adjustment has the largest effect on the most recent data years for cancers that are frequently diagnosed in outpatient settings (eg, melanoma, leukemia, and prostate cancer) and provides the most accurate portrayal of cancer occurrence in the most recent time period.20 For example, the leukemia incidence rate for 2015 in the 9 oldest SEER registries was 12% higher after adjusting for reporting delays (15.2 vs 13.6 per 100,000 population).10

### Projected Cancer Cases and Deaths in 2019

The most recent year for which reported incidence and mortality data are available lags 2 to 4 years behind the current year due to the time required for data collection, compilation, quality control, and dissemination. Therefore, we projected the numbers of new cancer cases and deaths in the United States in 2019 to provide an estimate of the contemporary cancer burden.

To calculate the number of invasive cancer cases, a generalized linear mixed model was used to estimate complete counts for each county (or health service area for rare cancers) from 2001 through 2015 using delay-adjusted, high-quality incidence data from 48 states and the District of Columbia (96% population coverage) and geographic variations in sociodemographic and lifestyle factors, medical settings, and cancer screening behaviors.21 (Data were unavailable for all years for Kansas and Minnesota, as well as for a few sporadic years for a handful of states.) Modeled counts were aggregated to the national and state level for each year, and a time series projection method (vector autoregression) was applied to all 15 years to estimate cases for 2019. Basal cell and squamous cell skin cancers cannot be estimated because incidence data are not collected by most cancer registries. For complete details of the case projection methodology, please refer to Zhu et al.22

New cases of in situ female breast carcinoma and melanoma of the skin diagnosed in 2019 were estimated by first approximating the number of cases occurring annually from 2006 through 2015 based on age-specific NAACCR incidence rates (data from 46 states with high-quality data for all 10 years) and US Census Bureau population estimates obtained via SEER\*Stat. Counts were then adjusted for delays in reporting using SEER delay factors for invasive disease (delay factors are unavailable for in situ cases) and projected to 2019 based on the average annual percent change generated by the joinpoint regression model.

The number of cancer deaths expected to occur in 2019 was estimated based on the most recent joinpoint-generated annual percent change in reported cancer deaths from 2002 through 2016 at the state and national levels as reported to the NCHS. For the complete details of this methodology, please refer to Chen et al.23

### **Other Statistics**

The number of cancer deaths averted in men and women due to the reduction in cancer death rates since the early 1990s was estimated by summing the difference between the annual number of recorded cancer deaths from the number that would have been expected if cancer death rates had remained at their peak. The expected number of deaths was estimated by applying the 5-year age- and sex-specific cancer death rates in the peak year for age-standardized cancer death rates (1990 in men and 1991 in women) to the corresponding age- and sex-specific populations in subsequent years through 2016.

Temporal trends in socioeconomic disparities in cancer mortality were examined using county-level poverty as a proxy for socioeconomic status. Cancer death rates by county-level poverty quintile were calculated using linked attributes from the US Census Bureau American Community Survey 2012–2016 available through SEER\*Stat. The total resident population in each quintile was 73,559,180 persons (1.81%-10.84% poverty); 62,695,449 persons (10.85%-14.10% poverty); 74,157,401 persons (14.11%-17.16% poverty); 76,945,467 persons (17.17%-21.17% poverty); and 35,770,016 persons (21.18%-53.95% poverty), respectively. County-level poverty in the United States has shifted slightly from the South to the West since 1970, although the highest concentration remains in the South.24

# **Selected Findings**

## Expected Numbers of New Cancer Cases

Table 1 presents the estimated numbers of new cases of invasive cancer in the United States in 2019 by sex and cancer type. In total, there will be approximately 1,762,450 cancer cases diagnosed, which is the equivalent of more than 4,800 new cases each day. In addition, there will be approximately 62,930 new cases of female breast carcinoma in situ and 95,830 new cases of melanoma in situ of the skin. The estimated numbers of new cases by state are shown in Table 2.

 Table 1. Estimated New Cancer Cases and Deaths by Sex, United States, 2019<sup>a</sup>

	ESTIMAT	ED NEW C	ASES	ESTIMATED DEATHS			
	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	FEMALE	
All sites	1,762,450	870,970	891,480	606,880	321,670	285,210	
Oral cavity & pharynx	53,000	38,140	14,860	10,860	7,970	2,890	
Tongue	17,060	12,550	4,510	3,020	2,220	800	
Mouth	14,310	8,430	5,880	2,740	1,800	940	
Pharynx	17,870	14,450	3,420	3,450	2,660	790	
Other oral cavity	3,760	2,710	1,050	1,650	1,290	360	
Digestive system	328,030	186,080	141,950	165,460	97,110	68,350	
Esophagus	17,650	13,750	3,900	16,080	13,020	3,060	
Stomach	27,510	17,230	10,280	11,140	6,800	4,340	
Small intestine	10,590	5,610	4,980	1,590	890	700	
Colon <sup>a</sup>	101,420	51,690	49,730	51,020	27,640	23,380	
Rectum	44,180	26,810	17,370				
Anus, anal canal, & anorectum	8,300	2,770	5,530	1,280	520	760	
Liver & intrahepatic bile duct	42,030	29,480	12,550	31,780	21,600	10,180	
Gallbladder & other biliary	12,360	5,810	6,550	3,960	1,610	2,350	
Pancreas	56,770	29,940	26,830	45,750	23,800	21,950	

Eye & orbit	3,360	1,860	1,500	370	200	170
Ureter & other urinary organs	3,930	2,630	1,300	980	600	380
Kidney & renal pelvis	73,820	44,120	29,700	14,770	9,820	4,950
Urinary bladder	80,470	61,700	18,770	17,670	12,870	4,800
Urinary system	158,220	108,450	49,770	33,420	23,290	10,130
Penis & other genital, male	2,080	2,080		410	410	
Testis	9,560	9,560		410	410	
Prostate	174,650	174,650		31,620	31,620	
Vagina & other genital, female	5,350		5,350	1,430		1,430
Vulva	6,070		6,070	1,280		1,280
Ovary	22,530		22,530	13,980		13,980
Uterine corpus	61,880		61,880	12,160		12,160
Uterine cervix	13,170		13,170	4,250		4,250
Genital system	295,290	186,290	109,000	65,540	32,440	33,100
Breast	271,270	2,670	268,600	42,260	500	41,760
Other nonepithelial skin	7,870	5,100	2,770	4,420	3,290	1,130
Melanoma of the skin	96,480	57,220	39,260	7,230	4,740	2,490
squamous)						
Skin (excluding basal &	104,350	62,320	42,030	11,650	8,030	3,620
Soft tissue (including heart)	12,750	7,240	5,510	5,270	2,840	2,430
Bones & joints	3,500	2,030	1,470	1,660	960	700
Other respiratory organs	5,880	4,070	1,810	1,080	720	360
Lung & bronchus	228,150	116,440	111,710	142,670	76,650	66,020
Larynx	12,410	9,860	2,550	3,760	3,010	750
Respiratory system	246,440	130,370	116,070	147,510	80,380	67,130
Other digestive organs	7,220	2,990	4,230	2,860	1,230	1,630

Brain & other nervous system	23,820	13,410	10,410	17,760	9,910	7,850
Endocrine system	54,740	15,650	39,090	3,210	1,560	1,650
Thyroid	52,070	14,260	37,810	2,170	1,020	1,150
Other endocrine	2,670	1,390	1,280	1,040	540	500
Lymphoma	82,310	45,660	36,650	20,970	12,100	8,870
Hodgkin lymphoma	8,110	4,570	3,540	1,000	590	410
Non-Hodgkin lymphoma	74,200	41,090	33,110	19,970	11,510	8,460
Myeloma	32,110	18,130	13,980	12,960	6,990	5,970
Leukemia	61,780	35,920	25,860	22,840	13,150	9,690
<b>Leukemia</b> Acute lymphocytic leukemia	<b>61,780</b> 5,930	<b>35,920</b> 3,280	<b>25,860</b> 2,650	<b>22,840</b> 1,500	<b>13,150</b> 850	<b>9,690</b> 650
<b>Leukemia</b> Acute lymphocytic leukemia Chronic lymphocytic leukemia	<b>61,780</b> 5,930 20,720	<b>35,920</b> 3,280 12,880	<b>25,860</b> 2,650 7,840	<b>22,840</b> 1,500 3,930	<b>13,150</b> 850 2,220	<b>9,690</b> 650 1,710
Leukemia         Acute lymphocytic leukemia         Chronic lymphocytic leukemia         Acute myeloid leukemia	<b>61,780</b> 5,930 20,720 21,450	<b>35,920</b> 3,280 12,880 11,650	<b>25,860</b> 2,650 7,840 9,800	<b>22,840</b> 1,500 3,930 10,920	<b>13,150</b> 850 2,220 6,290	<b>9,690</b> 650 1,710 4,630
Leukemia   Acute lymphocytic leukemia   Chronic lymphocytic leukemia   Acute myeloid leukemia   Chronic myeloid leukemia	<b>61,780</b> 5,930 20,720 21,450 8,990	<b>35,920</b> 3,280 12,880 11,650 5,250	<b>25,860</b> 2,650 7,840 9,800 3,740	<b>22,840</b> 1,500 3,930 10,920 1,140	<b>13,150</b> 850 2,220 6,290 660	<b>9,690</b> 650 1,710 4,630 480
Leukemia   Acute lymphocytic leukemia   Chronic lymphocytic leukemia   Acute myeloid leukemia   Chronic myeloid leukemia   Other leukemia <sup>a</sup>	<b>61,780</b> 5,930 20,720 21,450 8,990 4,690	<b>35,920</b> 3,280 12,880 11,650 5,250 2,860	<b>25,860</b> 2,650 7,840 9,800 3,740 1,830	22,840 1,500 3,930 10,920 1,140 5,350	<b>13,150</b> 850 2,220 6,290 660 3,130	<b>9,690</b> 650 1,710 4,630 480 2,220

\* Rounded to the nearest 10; cases exclude basal cell and squamous cell skin cancers and in situ carcinoma except urinary bladder. Approximately 62,930 cases of carcinoma in situ of the female breast and 95,830 cases of melanoma in situ will be newly diagnosed in 2019.

† Deaths for colon and rectal cancers are combined because a large number of deaths from rectal cancer are misclassified as colon.

<sup>‡</sup> More deaths than cases may reflect a lack of specificity in recording the underlying cause of death on death certificates and/or an undercount in the case estimate.

Note: These are model-based estimates that should be interpreted with caution and not compared with those for previous years.

STATE	ALL CASES	FEMALE BREAST	UTERINE CERVIX	COLON & RECTUM	UTERINE CORPUS	LEUKEMIA	LUNG & BRONCHUS	ME
Alabama	28,950	4,240	240	2,330	760	840	4,150	
Alaska	3,090	470	а	290	110	90	400	
Arizona	37,490	5,630	250	2,840	1,200	1,110	4,290	
Arkansas	16,580	2,210	140	1,440	510	560	2,690	
California	186,920	27,700	1,590	15,360	6,230	6,030	18,990	
Colorado	26,800	4,180	170	1,940	830	810	2,690	
Connecticut	21,950	3,490	120	1,560	720	670	2,580	
Delaware	5,870	930	а	440	220	210	840	
Dist. of Columbia	3,190	510	а	260	120	80	340	
Florida	131,470	19,130	1,040	11,310	4,520	4,980	18,560	
Georgia	50,450	8,000	440	4,450	1,640	1,800	7,070	
Hawaii	7,120	1,280	50	620	310	200	860	
Idaho	8,390	1,340	50	630	310	340	1,030	
Illinois	68,560	11,560	510	6,030	2,700	2,380	9,130	
Indiana	35,280	5,820	270	3,360	1,330	1,230	5,500	
lowa	17,810	2,730	100	1,540	660	730	2,410	
Kansas	15,340	2,420	110	1,290	520	590	2,000	
Kentucky	26,400	3,670	200	2,320	890	940	4,960	
Louisiana	26,800	3,770	230	2,340	700	830	3,810	
Maine	8,920	1,390	50	670	320	310	1,400	
Maryland	33,140	5,290	230	2,620	1,280	960	4,040	

Massachusetts	40,020	6,610	210	2,840	1,380	1,140	5,150
Michigan	58,360	9,310	360	5,040	2,200	1,930	8,070
Minnesota	30,560	4,740	140	2,300	1,080	1,360	3,600
Mississippi	17,050	2,370	150	1,680	450	520	2,520
Missouri	35,480	5,350	260	3,110	1,180	1,240	5,490
Montana	5,920	890	а	470	220	240	820
Nebraska	9,780	1,580	70	900	360	420	1,290
Nevada	14,810	2,190	140	1,340	420	530	1,880
New	8,610	1,330	а	590	300	260	1,140
Hampshire							
New Jersey	53,400	8,340	410	4,250	2,130	2,070	6,070
New Mexico	9,460	1,440	80	830	370	360	1,070
New York	111,870	17,490	880	9,150	4,500	4,540	13,380
North Carolina	58,690	8,870	410	4,310	1,960	1,960	8,010
North Dakota	3,940	590	а	350	130	170	430
Ohio	67,150	10,240	430	6,200	2,600	2,100	9,680
Oklahoma	20,540	2,980	170	1,840	630	780	3,220
Oregon	23,320	3,390	150	1,620	810	670	2,900
Pennsylvania	79,890	12,070	540	6,520	3,280	3,040	10,380
Rhode Island	6,540	1,010	а	470	210	190	940
South Carolina	29,830	4,470	210	2,370	930	1,040	4,360
South Dakota	4,770	750	а	430	160	200	580
Tennessee	37,350	5,580	310	3,290	1,210	1,280	6,210
Texas	124,890	18,750	1,290	10,950	4,090	4,820	14,750
Utah	11,620	1,660	70	770	420	480	780
Vermont	3,920	620		280	130	130	510

United States	1,762,450	268,600	13,170	145,600	61,880	61,780	228,150	
Wyoming	2,930	440	а	250	100	110	310	
Wisconsin	34,220	5,270	190	2,450	1,290	1,320	4,150	
West Virginia	12,440	1,540	80	980	450	410	2,010	
Washington	39,160	5,840	230	2,800	1,400	1,370	4,770	
Virginia	45,440	7,120	310	3,540	1,650	1,400	5,950	
			а					

\* Rounded to the nearest 10; excludes basal cell and squamous cell skin cancers and in situ carcinomas except urinary bladder. Estimates for Puerto Rico are not available.

† Estimate is fewer than 50 cases.

Note: These are model-based estimates that should be interpreted with caution and not compared with those for previous years. State estimates may not add to US total due to rounding and the exclusion of states with fewer than 50 cases.

Figure 1 depicts the most common cancers expected to be diagnosed in men and women in 2019. Prostate, lung and bronchus (referred to as lung hereafter), and colorectal cancers (CRCs) account for 42% of all cases in men, with prostate cancer alone accounting for nearly 1 in 5 new diagnoses. For women, the 3 most common cancers are breast, lung, and colorectum, which collectively represent one-half of all new diagnoses; breast cancer alone accounts for 30% of all new cancer diagnoses in women.

			Males	Femal	es		
Prostate	174,650	20%		_	Breast	268,600	30%
Lung & bronchus	116,440	13%			Lung & bronchus	111,710	13%
Colon & rectum	78,500	9%		T	Colon & rectum	67,100	8%
Urinary bladder	61,700	7%			Uterine corpus	61,880	7%
Melanoma of the skin	57,220	7%			Melanoma of the skin	39,260	4%
Kidney & renal pelvis	44,120	5%			Thyroid	37,810	4%
Non-Hodgkin lymphoma	41,090	5%			Non-Hodgkin lymphoma	33,110	4%
Oral cavity & pharynx	38,140	4%			Kidney & renal pelvis	29,700	3%
Leukemia	35,920	4%			Pancreas	26,830	3%
Pancreas	29,940	3%			Leukemia	25,860	3%
All Sites	870 970	100%			All Citor	891 480	100%
mated Deaths	010,010	100 /8	_		All offes		100 /
mated Deaths			Males	Femal	All Sites		10070
mated Deaths	76,650	24%	Males	Femal	es Lung & bronchus	66,020	23%
Lung & bronchus Prostate	76,650 31,620	24% 10%	Males	Femal	es Lung & bronchus Breast	66,020 41,760	23%
Lung & bronchus Prostate Colon & rectum	76,650 31,620 27,640	24% 10% 9%	Males	Femal	es Lung & bronchus Breast Colon & rectum	66,020 41,760 23,380	23% 15% 8%
Imated Deaths Lung & bronchus Prostate Colon & rectum Pancreas	76,650 31,620 27,640 23,800	24% 10% 9% 7%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas	66,020 41,760 23,380 21,950	23% 15% 8%
mated Deaths Lung & bronchus Prostate Colon & rectum Pancreas Liver & intrahepatic bile duct	76,650 31,620 27,640 23,800 21,600	24% 10% 9% 7% 7%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas Ovary	66,020 41,760 23,380 21,950 13,980	23% 15% 8% 5%
mated Deaths Lung & bronchus Prostate Colon & rectum Pancreas Liver & intrahepatic bile duct Leukemia	76,650 31,620 27,640 23,800 21,600 13,150	24% 10% 9% 7% 7% 4%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas Ovary Uterine corpus	66,020 41,760 23,380 21,950 13,980 12,160	23% 15% 8% 5% 4%
mated Deaths Lung & bronchus Prostate Colon & rectum Pancreas Liver & intrahepatic bile duct Leukemia Esophagus	76,650 31,620 27,640 23,800 21,600 13,150 13,020	24% 10% 9% 7% 4% 4%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas Ovary Uterine corpus Liver & intrahepatic bile duct	66,020 41,760 23,380 21,950 13,980 12,160 10,180	23% 15% 8% 5% 4% 4%
mated Deaths Lung & bronchus Prostate Colon & rectum Pancreas Liver & intrahepatic bile duct Leukemia Esophagus Urinary bladder	76,650 31,620 27,640 23,800 21,600 13,150 13,020 12,870	24% 10% 9% 7% 4% 4%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas Ovary Uterine corpus Liver & intrahepatic bile duct Leukemia	66,020 41,760 23,380 21,950 13,980 12,160 10,180 9,690	23% 15% 8% 5% 4% 3%
mated Deaths Lung & bronchus Prostate Colon & rectum Pancreas Liver & intrahepatic bile duct Leukemia Esophagus Urinary bladder Non-Hodgkin lymphoma	76,650 31,620 27,640 23,800 21,600 13,150 13,020 12,870 11,510	24% 10% 9% 7% 4% 4% 4%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas Ovary Uterine corpus Liver & intrahepatic bile duct Leukemia Non-Hodgkin lymphoma	66,020 41,760 23,380 21,950 13,980 12,160 10,180 9,690 8,460	23% 15% 8% 5% 4% 3% 3%
imated Deaths Lung & bronchus Prostate Colon & rectum Pancreas Liver & intrahepatic bile duct Leukemia Esophagus Urinary bladder Non-Hodgkin lymphoma Brain & other nervous system	76,650 31,620 27,640 23,800 21,600 13,150 13,020 12,870 11,510 9,910	24% 10% 9% 7% 4% 4% 4% 4% 3%	Males	Femal	es Lung & bronchus Breast Colon & rectum Pancreas Ovary Uterine corpus Liver & intrahepatic bile duct Leukemia Non-Hodgkin lymphoma Brain & other nervous system	66,020 41,760 23,380 21,950 13,980 12,160 10,180 9,690 8,460 7,850	23% 15% 8% 5% 4% 3% 3% 3%

#### Open in figure viewer | PowerPoint

Ten Leading Cancer Types for the Estimated New Cancer Cases and Deaths by Sex, United States, 2019. Estimates are rounded to the nearest 10 and exclude basal cell and squamous cell skin cancers and in situ carcinoma except urinary bladder. Ranking is based on modeled projections and may differ from the most recent observed data.

The lifetime probability of being diagnosed with invasive cancer is slightly higher for men (39.3%) than for women (37.7%) (Table 3). The reasons for the excess risk in men are not fully understood, but partly reflect differences in environmental exposures, endogenous hormones, and probably complex interactions between these influences. Recent research suggests that sex differences in immune function and response may also play a role.25 Adult height, which is determined by genetics and childhood nutrition, is positively associated with cancer incidence and mortality in both men and women,26 and has been estimated to account for one-third of the sex disparity.27

Table 3. Probability (%) of Developing Invasive Cancer Within Selected Age Intervals by Sex,

### United States, 2013 to 2015<sup>a</sup>

		BIRTH TO 49	50 TO 59	60 TO 69	≥70	BIRTH TO DEATH
All sites <sup>†</sup>	Male	3.4 (1 in 30)	6.1 (1 in 16)	13.2 (1 in 8)	31.9 (1 in 3)	39.3 (1 in 3)
	Female	5.6 (1 in 18)	6.2 (1 in 16)	10.0 (1 in 10)	26.0 (1 in 4)	37.7 (1 in 3)
Breast	Female	2.0 (1 in 51)	2.3 (1 in 43)	3.5 (1 in 29)	6.7 (1 in 15)	12.4 (1 in 8)
Colorectum	Male	0.4 (1 in 272)	0.7 (1 in 143)	1.2 (1 in 87)	3.3 (1 in 30)	4.4 (1 in 23)
	Female	0.3 (1 in 292)	0.5 (1 in 190)	0.8 (1 in 123)	3.0 (1 in 33)	4.1 (1 in 25)
Kidney & renal pelvis	Male	0.2 (1 in 440)	0.4 (1 in 280)	0.6 (1 in 155)	1.3 (1 in 73)	2.1 (1 in 47)
	Female	0.2 (1 in 665)	0.2 (1 in 575)	0.3 (1 in 319)	0.7 (1 in 135)	1.2 (1 in 82)
Leukemia	Male	0.3 (1 in 396)	0.2 (1 in 570)	0.4 (1 in 259)	1.4 (1 in 72)	1.8 (1 in 56)
	Female	0.2 (1 in 508)	0.1 (1 in 876)	0.2 (1 in 434)	0.9 (1 in 112)	1.3 (1 in 80)
Lung & bronchus	Male	0.1 (1 in 719)	0.6 (1 in 158)	1.8 (1 in 56)	6.0 (1 in 16)	6.7 (1 in 15)
	Female	0.1 (1 in 673)	0.6 (1 in 178)	1.4 (1 in 72)	4.7 (1 in 21)	5.9 (1 in 17)
Melanoma of the skin ‡	Male	0.5 (1 in 215)	0.5 (1 in 186)	1.0 (1 in 104)	2.7 (1 in 37)	3.7 (1 in 27)
	Female	0.7 (1 in 150)	0.4 (1 in 238)	0.5 (1 in 191)	1.1 (1 in 87)	2.5 (1 in 40)

Non-Hodgkin	Male	0.3 (1 in	0.3 (1 in	0.6 (1 in	1.8 (1 in	2.4 (1 in 42)
lymphoma		382)	350)	176)	54)	
	Female	0.2 (1 in	0.2 (1 in	0.4 (1 in	1.4 (1 in	1.9 (1 in 54)
		548)	484)	247)	74)	
Prostate	Male	0.2 (1 in	1.7 (1 in	4.6 (1 in	7.9 (1 in	11.2 (1 in 9)
		437)	59)	22)	13)	
Thyroid	Male	0.2 (1 in	0.1 (1 in	0.2 (1 in	0.2 (1 in	0.6 (1 in 156)
		513)	764)	584)	417)	
	Female	0.8 (1 in	0.4 (1 in	0.3 (1 in	0.4 (1 in	1.8 (1 in 55)
		122)	268)	286)	262)	
Uterine cervix	Female	0.3 (1 in	0.1 (1 in	0.1 (1 in	0.2 (1 in	0.6 (1 in 162)
		366)	835)	938)	628)	
Uterine corpus	Female	0.3 (1 in	0.6 (1 in	1.0 (1 in	1.3 (1 in	2.9 (1 in 35)
		333)	164)	102)	75)	

\* For people without a history of cancer at beginning of age interval.

† All sites excludes basal cell and squamous cell skin cancers and in situ cancers except urinary bladder.

**‡** Probabilities for non-Hispanic whites only.

### Expected Number of Cancer Deaths

An estimated 606,880 Americans will die from cancer in 2019, corresponding to almost 1,700 deaths per day (Table 1). The greatest number of deaths are from cancers of the lung, prostate, and colorectum in men and the lung, breast, and colorectum in women (Fig. 1). One-quarter of all cancer deaths are due to lung cancer. Table 4 provides the estimated numbers of cancer deaths in 2019 by state.

 Table 4. Estimated Deaths for Selected Cancers by State, 2019<sup>3</sup>

STATE	ALL	BRAIN &	FEMALE	COLON	LEUKEMIA	LIVER &	LUNG &
	SITES	OTHER	BREAST	&		INTRAHEPATIC	BRONCHUS
		NERVOUS		RECTUM		BILE DUCT	

		SYSTEM					
Alabama	10,630	350	690	930	380	540	2,760
Alaska	1,120	а	70	110	а	60	260
Arizona	12,470	400	890	1,050	510	710	2,630
Arkansas	6,800	190	410	600	240	310	1,960
California	60,590	1,970	4,560	5,290	2,400	4,070	10,970
Colorado	8,120	290	610	660	330	430	1,500
Connecticut	6,470	210	430	470	270	320	1,440
Delaware	2,140	60	150	150	80	110	540
Dist. of Columbia	1,020	а	100	100	а	90	180
Florida	45,000	1,240	3,000	3,700	1,740	2,300	10,880
Georgia	17,880	530	1,350	1,630	590	940	4,340
Hawaii	2,560	50	160	230	80	190	550
Idaho	3,040	110	220	250	110	160	620
Illinois	24,410	670	1,720	2,070	900	1,150	5,940
Indiana	13,690	360	870	1,110	510	580	3,690
lowa	6,480	200	380	560	240	270	1,600
Kansas	5,550	170	350	470	240	260	1,370
Kentucky	10,580	290	610	820	370	460	3,290
Louisiana	9,260	230	620	830	320	580	2,390
Maine	3,310	100	180	230	110	120	890
Maryland	10,780	300	830	880	390	600	2,380
Massachusetts	12,420	400	750	870	480	690	2,920
Michigan	21,150	600	1,410	1,650	770	920	5,410

Minnesota	10,020	320	640	790	420	440	2,260
Mississippi	6,720	190	440	650	210	340	1,810
Missouri	13,080	340	860	1,050	480	580	3,650
Montana	2,100	70	140	180	80	100	480
Nebraska	3,520	120	230	310	150	130	840
Nevada	5,390	200	400	540	200	250	1,280
New Hampshire	2,820	90	180	200	100	120	730
New Jersey	15,860	470	1,250	1,410	590	750	3,390
New Mexico	3,720	100	270	340	130	250	700
New York	35,010	940	2,460	2,890	1,370	1,740	7,790
North Carolina	20,410	550	1,390	1,580	720	1,110	5,370
North Dakota	1,280	а	80	120	50	а	300
Ohio	25,440	680	1,710	2,110	920	1,100	6,690
Oklahoma	8,420	220	540	760	340	420	2,270
Oregon	8,270	250	560	650	300	500	1,820
Pennsylvania	28,170	770	1,900	2,380	1,080	1,320	6,730
Rhode Island	2,140	60	130	160	80	120	560
South Carolina	10,720	300	740	870	380	530	2,710
South Dakota	1,680	60	110	170	70	70	410
Tennessee	14,840	360	950	1,220	520	730	4,190
Texas	41,300	1,300	2,980	3,850	1,580	2,810	8,640
Utah	3,310	140	280	280	160	170	440
Vermont	1,440	50	80	110	50	50	370
Virginia	15,200	440	1,120	1,340	520	770	3,590
Washington	13,010	430	890	1,000	480	730	2,830

United States	606,880	17,760	41,760	51,020	22,840	31,780	142,670
Wyoming	980	а	70	80	50	60	200
Wisconsin	11,730	380	740	900	490	480	2,770
West Virginia	4,820	120	290	440	190	190	1,360

\* Rounded to the nearest 10. Estimates for Puerto Rico are not available.

† Estimate is fewer than 50 deaths.

Note: These are model-based estimates that should be interpreted with caution and not compared with those for previous years. State estimates may not add to US total due to rounding and the exclusion of states with fewer than 50 deaths.

### Trends in Cancer Incidence

Figure 2 illustrates long-term trends in cancer incidence rates for all cancers combined by sex. Cancer incidence patterns reflect trends in behaviors associated with cancer risk and changes in medical practice, such as the use of cancer screening tests. The volatility in incidence for males reflects rapid changes in prostate cancer incidence rates, which spiked in the late 1980s and early 1990s (Fig. 3) due to a surge in the detection of asymptomatic disease as a result of widespread prostate-specific antigen (PSA) testing among previously unscreened men.28



#### Open in figure viewer PowerPoint

Trends in Cancer Incidence (1975 to 2015) and Mortality Rates (1975 to 2016) by Sex, United States. Rates are age adjusted to the 2000 US standard population. Incidence rates also are adjusted for delays in reporting.



#### Open in figure viewer | PowerPoint

Trends in Incidence Rates for Selected Cancers by Sex, United States, 1975 to 2015. Rates are age adjusted to the 2000 US standard population and adjusted for delays in reporting. \*Includes intrahepatic bile duct.

Over the past decade of data, the overall cancer incidence rate in men declined by approximately 2% per year (Table 5). This trend reflects accelerated declines during the past 5 data years (2011-2015) of approximately 3% per year for cancers of the lung and colorectum, and 7% per year for prostate cancer. The sharp drop in prostate cancer incidence has been attributed to decreased PSA testing from 2008 to 2013 in the wake of US Preventive Services Task Force recommendations against the routine use of the test to screen for prostate cancer (Grade D) in men aged 75 years and older in 2008 and in all men in 2011 because of growing concerns about overdiagnosis and overtreatment.29, 30 Although PSA testing prevalence stabilized from 2013 to 2015,31 the effect of the reduction in screening on the occurrence of advanced disease is being watched closely. Based on analysis of cancer registry data covering 89% of the US population, Negoita et al recently reported that the overall decline in prostate cancer incidence masks an increase in distant stage diagnoses since around 2010 across age and race, although improved staging may have contributed to this trend.32 The Task Force has revised their recommendation for men aged 55 to 69 years to informed decision making (Grade C) based on an updated evidence review, noting that "screening offers a small potential benefit" of reduced prostate cancer

mortality "in some men."33-35

**Table 5.** Trends in Delay-Adjusted Incidence Rates for Selected Cancers by Sex, UnitedStates, 1975 to 2015

	TREND	1	TREND	2	TREND	3	TREND	4	TREND	5	TRE
	YEARS	APC	YEARS	APC	YEARS	APC	YEARS	APC	YEARS	APC	YEA
All sites	·			<u>.</u>		·		<u>.</u>		<u>.</u>	-
Overall	1975- 1989	1.2 <sup>8</sup>	1989- 1992	2.8	1992- 1995	-2.4	1995- 1998	1.1	1998- 2009	- 0.3 <sup>2</sup>	200 201
Male	1975- 1989	1.3 <sup>a</sup>	1989- 1992	5.2 <sup>a</sup>	1992- 1995	-4.9 <sup>a</sup>	1995- 1999	0.6	1999- 2008	- 0.6 <sup>2</sup>	200 201
Female	1975- 1979	-0.3	1979- 1987	1.6 <sup>a</sup>	1987- 1995	0.1	1995- 1998	1.5	1998- 2003	-0.6	200 201
Female breast	1975- 1980	-0.5	1980- 1987	4.0 <sup>a</sup>	1987- 1994	-0.2	1994- 1999	1.8 <sup>a</sup>	1999- 2004	- 2.3 <sup>a</sup>	200 201
Colorectum											
Male	1975- 1985	1.1 <sup>a</sup>	1985- 1991	-1.2 <sup>a</sup>	1991- 1995	-3.2 <sup>a</sup>	1995- 1998	2.1	1998- 2015	- 2.9 <sup>a</sup>	
Female	1975- 1985	0.3	1985- 1995	-1.9 <sup>a</sup>	1995- 1998	1.8	1998- 2008	- 2.0 <sup>2</sup>	2008- 2011	- 4.6 <sup>a</sup>	201 20 <sup>7</sup>
Liver & intrahepatic bile duct											
Male	1975- 1984	2.2 <sup>a</sup>	1984- 2011	3.9 <sup>3</sup>	2011- 2015	1.0					
Female	1975- 1983	0.4	1983- 1998	4.4 <sup>a</sup>	1998- 2001	-0.4	2001- 2015	3.4 <sup>a</sup>			

Lung &

bronchus											
Male	1975-	1.5 <sup>a</sup>	1982-	-0.5 <sup>2</sup>	1991-	-1.7	2008-	-			
	1982		1991		2008		2015	2.9 <sup>a</sup>			
Female	1975-	5.6 <sup>a</sup>	1982-	3.4	1991-	0.5 <sup>a</sup>	2006-	-			
	1982		1991		2006		2015	1.5 <sup>a</sup>			
Melanoma											
of skin											
Male	1975-	5.4 <sup>a</sup>	1986-	3.1 <sup>a</sup>	2005-	1.8 <sup>2</sup>					
	1986		2005		2015						
Female	1975-	5.5 <sup>a</sup>	1980-	2.4	2009-	-1.3	2012 <b>-</b>	5.4 <sup>a</sup>			
	1980		2009		2012		2015				
Pancreas											
Male	1975-	-1.8 <sup>a</sup>	1981-	1.2	1985-	-2.2 <sup>2</sup>	1990-	0.2	2003-	3.1	200
	1981		1985		1990		2003		2006		20
Female	1975-	1.4 <sup>a</sup>	1984-	-0.5	1996-	1.0 <sup>a</sup>					
	1984		1996		2015						
Prostate	1975-	2.6 <sup>a</sup>	1988-	16.5 <sup>2</sup>	1992-	-	1995-	2.2	2000-	-	200
	1988		1992		1995	11.5 <sup>a</sup>	2000		2009	1.6 <sup>a</sup>	20
Thyroid											
Male	1975-	-4.6	1980-	1.8	1997-	5.5 <sup>a</sup>	2012-	-1.1			
	1980		1997		2012		2015				
Female	1975-	6.5	1977-	-5.2	1980-	2.3	1993-	4.4 <sup>a</sup>	1999-	7.1 <sup>a</sup>	200
	1977		1980		1993		1999		2009		20
Uterine	1975-	-6.0 <sup>2</sup>	1979-	-1.7 <sup>a</sup>	1988-	0.7	1997-	-	2006-	3.5 <sup>a</sup>	200
corpus	1979		1988		1997		2006	0.4	2009		20

AAPC indicates average annual percent change; APC, annual percent change based on delay-adjusted incidence rates age adjusted to the 2000 US standard population.

Note: Trends analyzed by the Joinpoint Regression Program, version 4.6, allowing up to 5 joinpoints. Trends are based on Surveillance, Epidemiology, and End Results (SEER) 9 areas. \* The APC or AAPC is significantly different from zero (P < .05).

The overall cancer incidence rate in women has remained generally stable over the past few decades. Declines have continued for lung cancer, but tapered in recent years for CRC, whereas rates for other common cancers are increasing or stable (Table 5). Breast cancer incidence rates increased from 2006 to 2015 by approximately 0.3% to 0.4% per year among non-Hispanic white (NHW) and Hispanic women, by 0.7% to 0.8% per year among black (non-Hispanic) and American Indian/Alaska Native women, and by 1.8% per year among Asian/Pacific Islander women.36 This trend may in part be a consequence of the obesity epidemic, as well as declining parity.37, 38

Lung cancer incidence continues to decline twice as fast in men as in women, reflecting historical differences in tobacco uptake and cessation, as well as upturns in female smoking prevalence in some birth cohorts.39, 40 However, smoking patterns do not appear to explain the higher lung cancer incidence rates recently reported in young women compared with men born around the 1960s.41 In contrast, CRC incidence patterns are generally similar in men and women (Fig. 3), although in the past 5 data years rates have continued to decline by approximately 3% per year in men, but appear to have stabilized in women (Table 5). Reductions in CRC incidence prior to 2000 are attributed equally to changes in risk factors and the use of screening, which allows for the removal of premalignant lesions.42 However, more recent rapid declines are thought to primarily reflect the increased uptake of colonoscopy, which now is the predominant screening test.43, 44 Colonoscopy use among US adults aged 50 years and older tripled from 21% in 2000 to 60% in 2015.45 The rapid declines in overall CRC incidence rates mask an increase in adults aged younger than 55 years of almost 2% per year since the mid-1990s.7

Incidence rates continue to increase for melanoma and cancers of the liver, thyroid, uterine corpus, and pancreas. Liver cancer incidence is rising faster than that for any other cancer in both men and women.38 Notably, however, the majority (71%) of cases in the United States are potentially preventable because most risk factors are modifiable (eg, obesity, excess alcohol consumption, cigarette smoking, and hepatitis B and C viruses).46 Approximately 24% of cases are caused by chronic hepatitis C virus (HCV) infection, which confers the largest relative risk and is also the most common chronic blood-borne infection in the United States.47 Although there is exciting potential to avert much of the future burden of HCV-associated disease because of new, well-tolerated, antiviral therapies that achieve cure rates of greater than 90%,48 most infected individuals are undiagnosed. One-time screening has been recommended for baby boomers (those born between 1945 and 1965), who

account for three-fourths of affected individuals,49, 50 since 2012 and is now even mandated in several states.51 However, only 14% of the more than 76 million boomers reported having received HCV testing in 2015.52 Compounding the challenge is a 3-fold spike in acute HCV infections reported to the CDC from 2010 through 2016, after a decade of stable/declining rates, that is attributed to the opioid epidemic.53, 54 Fewer than 10% of new infections are reported and the CDC estimates the actual number of acute infections in 2016 to be 41,200 (95% confidence interval, 32,600-140,600), approximately 75% to 85% of which will progress to chronic infection.

### Cancer Survival

The 5-year relative survival rate for all cancers combined diagnosed during 2008 through 2014 was 67% in whites and 62% in blacks.10 Figure 4 shows 5-year relative survival rates by cancer type, stage at diagnosis, and race. For all stages combined, survival is highest for prostate cancer (98%), melanoma of the skin (92%), and female breast cancer (90%) and lowest for cancers of the pancreas (9%), liver (18%), esophagus (19%), and lung (19%). Black patients have lower survival rates than whites for every cancer type shown in Figure 4 except for cancers of the kidney and pancreas, with the absolute difference being 10% or higher for most. The largest disparities are for melanoma (26%) and cancers of the uterine corpus (21%) and oral cavity and pharynx (18%), in part reflecting a much later stage at diagnosis in black patients (Fig. 5). However, blacks also have lower stage-specific survival for most cancer types. After adjusting for sex, age, and stage at diagnosis, the relative risk of death after a cancer diagnosis is 33% higher in black patients than in white patients.55 The disparity is even larger for American Indians/Alaska Natives, who are 51% more likely than whites to die from their cancer.



#### Open in figure viewer PowerPoint

Five-Year Relative Survival Rates for Selected Cancers by Race and Stage at Diagnosis, United States, 2008 to 2014. \*The standard error of the survival rate is between 5 and 10 percentage points. †The survival rate for carcinoma in situ of the urinary bladder is 95% in all races, 95% in whites, and 91% in blacks.



#### Open in figure viewer | PowerPoint

Stage Distribution for Selected Cancers by Race, United States, 2008 to 2014. Stage categories do not sum to 100% because sufficient information was not available to stage all cases.

Cancer survival has improved since the mid-1970s for all of the most common cancers except those of the uterine cervix and uterine corpus,55 although for some cancer types (eg, breast and prostate) this partly reflects lead time bias because of changes in detection practice. Progress has been especially rapid for hematopoietic and lymphoid malignancies due to improvements in treatment protocols, including the discovery of targeted therapies. For example, the 5-year relative survival rate for chronic myeloid leukemia increased from 22% for patients diagnosed in the mid-1970s to 69% for those diagnosed during 2008 through 2014,10 and most patients treated with tyrosine kinase inhibitors experience nearly normal life expectancy.56

In contrast to the steady increase in survival for most cancer types, advances have been slow for lung and pancreatic cancers, partly because greater than one-half of cases are diagnosed at a distant stage (Fig. 5). There is a potential for earlier lung cancer diagnosis through screening with low-dose computed tomography, which has demonstrated a 20% reduction in lung cancer mortality in current/former smokers with a history of 30 or more packyears.57 However, the translation of this benefit from clinical trial participants to the general population remains challenging. In 2015, only 4% of the 6.8 million eligible Americans reported being screened for lung cancer with low-dose computed tomography.58 Another study found that more individuals who did not meet guideline-recommended criteria for lung cancer screening had received a recent test than those who did meet criteria.59 Broad implementation of guideline-recommended lung cancer screening will require new systems to facilitate unique aspects of the process, such as identifying eligible patients and acquainting physicians with information that should be delivered during the shared decision-making conversation, which is recommended by the American Cancer Society and US Preventive Services Task Force and required by the Centers for Medicare and Medicaid Services. A recent small study suggests stark failure in the practice of shared decision making by primary care and pulmonary physicians.60

### Trends in Cancer Mortality

Mortality rates are a better indicator of progress against cancer than incidence or survival rates because they are less affected by biases resulting from changes in detection practices.61 The cancer death rate rose during most of the 20th century, largely driven by rapid increases in lung cancer deaths among men as a consequence of the tobacco epidemic. However, since its peak of 215.1 deaths (per 100,000 population) in 1991, the cancer death rate has dropped steadily by approximately 1.5% per year, resulting in an overall decline of 27% as of 2016 (156.0 per 100,000 population). This translates to an estimated 2,629,200 fewer cancer deaths (1,804,000 in men and 825,200 in women) than what would have occurred if mortality rates had remained at their peak (Fig. 6). The number of averted deaths is larger for men than for women because the total decline in cancer mortality has been steeper for men (34% vs 24%).



#### Open in figure viewer PowerPoint

Total Number of Cancer Deaths Averted From 1991 to 2016 in Men and From 1992 to 2016 in Women, United States. The blue line represents the actual number of cancer deaths recorded in each year, and the red line represents the number of cancer deaths that would have been expected if cancer death rates had remained at their peak.

The decline in cancer mortality over the past 2 decades is primarily the result of steady reductions in smoking and advances in early detection and treatment, which are reflected in the rapid declines for the 4 major cancers (lung, breast, prostate, and colorectum) (Fig. 7). Specifically, the death rate for lung cancer dropped by 48% from 1990 to 2016 among males and by 23% from 2002 to 2016 among females, whereas the death rate for breast cancer dropped by 40% from 1989 to 2016, that for prostate cancer dropped by 51% from 1993 to 2016, and that for CRC dropped by 53% from 1970 to 2016. During the most recent data years, declines in mortality from lung cancer have accelerated whereas those for CRC have slowed (Table 6). Prostate cancer mortality stabilized during 2013 through 2016 after 2 decades of steep (4% per year) reductions that are attributed to an earlier stage at diagnosis due to PSA testing and advances in treatments.62, 63 The leveling of rates is temporally associated with both declines in PSA testing and an uptick in distant stage disease

diagnoses.32 Death rates rose from 2012 through 2016 for cancers of the liver, pancreas, and uterine corpus (Table 6), as well as for cancers of the brain and other nervous system, soft tissue (including heart), and sites within the oral cavity and pharynx associated with the human papillomavirus (HPV).1



#### Figure 7

#### Open in figure viewer | PowerPoint

Trends in Cancer Mortality Rates by Sex Overall and for Selected Cancers, United States, 1930 to 2016. Rates are age adjusted to the 2000 US standard population. Due to improvements in International Classification of Diseases (ICD) coding over time, numerator data for cancers of the lung and bronchus, colon and rectum, liver, and uterus differ from the contemporary time period. For example, rates for lung and bronchus include pleura, trachea, mediastinum, and other respiratory organs.

	TREN	D 1	TREN	D 2	TREN	D 3	TREN	D 4	TREN	D 5	TRENC
	YEARS	APC	YEARS								
All sites		·			·		·		·	<u> </u>	·
Overall	1975-	0.5 <sup>a</sup>	1984-	0.3	1991-	-0.5	1994-	-	1998-	-	2001-
	1984		1991		1994		1998	1.3 <sup>a</sup>	2001	0.8 <sup>a</sup>	2016
Male	1975-	1.0 <sup>a</sup>	1979-	0.3 <sup>2</sup>	1990-	-0.5	1993-	-	2001-	-	
	1979		1990		1993		2001	1.5 <sup>a</sup>	2016	1.8 <sup>a</sup>	
Female	1975-	0.6 <sup>a</sup>	1990-	-0.2	1994-	_	2002-	_			
	1990		1994		2002	0.8 <sup>a</sup>	2016	1.4 <sup>a</sup>			
Female	1975-	04	1990-	_	1995-	_	1998-	_			
breast	1990	0.1	1995	1.7	1998	3.4	2016	1.8			
Colorectum											
Male	1975-	0.6	1979-	_	1987-	_	2002-	-	2005-	-	
	1979		1987	0.0	2002	1.9 <sup>a</sup>	2005	4.2 <sup>a</sup>	2016	2.4*	
Female	1975-	-	1984-	_	2001-	_	2012-	_			
	1984	1.0 <sup>2</sup>	2001	1.8 <sup>a</sup>	2012	2.9 <sup>a</sup>	2016	1.5 <sup>a</sup>			
Liver &											
intrahepatic											
bile duct											
Male	1975-	1.5 <sup>a</sup>	1985-	3.8 <sup>2</sup>	1996-	0.3	1999-	2.7	2013-	0.7	
	1985		1996		1999		2013		2016		
Female	1975-	-1.5	1978-	1.4 <sup>a</sup>	1988-	3.9 <sup>a</sup>	1995-	0.4	2000-	1.6 <sup>a</sup>	2008-
	1978		1988		1995		2000		2008		2016
Lung &											
bronchus											
Male	1975-	2.4	1978-	1.2 <sup>a</sup>	1984-	0.3	1991-	-	2005-	-	2012-

	1978		1984		1991		2005	1.9 <sup>a</sup>	2012	2.9	2016
Female	1975-	5.9 <sup>a</sup>	1983-	3.8 <sup>a</sup>	1992-	0.5 <sup>a</sup>	2002-	-	2007-	-	2014-
	1983		1992		2002		2007	0.7 <sup>a</sup>	2014	2.0	2016
Melanoma											
of skin											
Male	1975-	2.3	1989-	0.3 <sup>2</sup>	2013-	-					
	1989		2013		2016	6.9 <sup>a</sup>					
Female	1975-	0.8	1988-	-	2014-	-					
	1988		2014	0.6	2016	9.3 <sup>2</sup>					
Pancreas											
Male	1975-	-	1986-	-	2000-	0.3					
	1986	0.8	2000	0.3	2016						
Female	1975-	0.8	1984-	0.1	2003-	1.0	2006-	0.0			
	1984		2003		2006		2016				
Prostate	1975-	0.9	1987-	3.0 <sup>2</sup>	1991-	-0.5	1994-	-	1998-	-	2013-
	1987		1991		1994		1998	4.2 <sup>a</sup>	2013	3.5 <sup>8</sup>	2016
Uterine	1975-	-	1993-	0.1	2008-	2.1 <sup>a</sup>					
corpus	1993	1.5 <sup>a</sup>	2008		2016						

AAPC indicates average annual percent change; APC, annual percent change based on mortality rates age adjusted to the 2000 US standard population.

Note: Trends analyzed by the Joinpoint Regression Program, version 4.6, allowing up to 5 joinpoints.

 $^{a}$  The APC or AAPC is significantly different from zero (P <.05).

### Recorded Number of Deaths in 2016

A total of 2,744,248 deaths were recorded in the United States in 2016, 22% of which were from cancer (Table 7). Cancer is the second leading cause of death after heart disease in both men and women nationally, but is the leading cause of death in many states,64 in Hispanic and Asian Americans,65, 66 and in people younger than 80 years. However, those 80 years and older are nearly 2 times more likely to die from heart disease than from

cancer. Among females, cancer is the first or second leading cause of death for every age group shown in Table 8, whereas among males, accidents, assault, and suicide predominate before age 40 years.

		_		_	_	_			_	
Tahlo 7	Ten	Ipading	Calicos	of Death	in the	Inited	States	2015	and 1	2016
	ICH	LCaung	Causes	U DCati		United	Juics,	2013		2010

		2015				2016		
		NO.	PERCENT	RATE	NO.	PERCENT	RATE	
RANK (2016)	All causes	2,712,630		733.0	2,744,248		729.1	-0.5%
1	Heart disease	633,842	23%	168.3	635,260	23%	165.5	-1.7%
2	Cancer	595,930	22%	158.7	598,038	22%	156.0	-1.7%
3	Accidents (unintentional injuries)	146,571	5%	43.1	161,374	6%	47.3	9.7%
4	Chronic lower respiratory diseases	155,041	6%	41.8	154,596	6%	40.8	-2.4%
5	Cerebrovascular disease	140,323	5%	37.6	142,142	5%	37.4	-0.5%
6	Alzheimer disease	110,561	4%	29.4	116,103	4%	30.3	3.1%
7	Diabetes mellitus	79,535	3%	21.3	80,058	3%	21.0	-1.4%
8	Influenza and pneumonia	57,062	2%	15.2	51,537	2%	13.6	-10.5%
9	Nephritis, nephrotic syndrome, & nephrosis	49,959	2%	13.4	50,046	2%	13.2	-1.5%

10	Intentional self-	44,193	2%	13.3	44,965	2%	13.4	0.8%
	harm (suicide)							

Death counts include unknown age.

Rates are per 100,000 population and age adjusted to the 2000 US standard population. Rank is based on number of deaths.

Source: National Center for Health Statistics, Centers for Disease Control and Prevention.

#### Table 8. Ten Leading Causes of Death in the United States by Age and Sex, 2016

	ALL	AGES	AGES 1	TO 19	AGES 2	0 TO 39
MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE
All Causes	All Causes	All Causes	All Causes	All Causes	All Causes	All Causes
1,400,232	1,344,016	13,110	7,250	79,366	35,290	230,142
1	Heart disease	Heart disease	Accidents (unintentional injuries)	Accidents (unintentional injuries)	Accidents (unintentional injuries)	Accidents (unintentional injuries)
	339,265	295,995	4,674	2,373	33,073	11,808
2	Cancer	Cancer	Assault (homicide)	Cancer	Intentional self-harm (suicide)	Cancer
	314,571	283,467	1,886	789	11,593	4,653
3	Accidents (unintentional injuries)	Cerebro- vascular disease	Intentional self-harm (suicide)	Intentional self-harm (suicide)	Assault (homicide)	Intentional self-harm (suicide)
	103,864	82,787	1,873	687	9,042	2,976
4	Chronic lower respiratory	Chronic lower respiratory	Cancer	Assault (homicide)	Heart disease	Heart disease

	diseases	diseases				
	73,045	81,551	1,064	555	5,362	2,621
5	Cerebro-	Alzheimer	Congenital	Congenital	Cancer	Assault
	vascular	disease	anomalies	anomalies		(homicide)
	disease					
		00 704	540	460	4.0.4.4	1 (10
	59,355	80,731	519	460	4,044	1,610
6	Diabetes	Accidents	Heart disease	Heart disease	Chronic liver	Pregnancy,
	mellitus	(unintentional			disease &	childbirth &
		injuries)			cirrhosis	puerperium
	43.763	57.510	341	258	1,255	864
	13,7 03	57,510		255	1,200	
7	Alzheimer	Diabetes	Chronic lower	Influenza &	Diabetes	Chronic liver
	disease	mellitus	respiratory	pneumonia	mellitus	disease &
			diseases			cirrhosis
	35,372	36,295	169	111	1,029	818
8	Intentional	Influenza &	Influenza &	Chronic lower	Cerebro-	Diabetes
	self-harm	pneumonia	pneumonia	respiratory	vascular	mellitus
	(suicide)			diseases	disease	
	34,727	26,526	133	105	784	722
9	Chronic liver	Nenhritis	Cerebro-	Senticemia	HIV disease	Cerebro-
	disease &	nephrotic	vascular			vascular
	cirrhosis	syndrome &	disease			disease
		nephrosis				
	25,818	24,647	120	87	727	576
10	Nephritis,	Septicemia	Septicemia	Cerebro-	Influenza &	Influenza &
	nephrotic			vascular	pneumonia	pneumonia
	syndrome &			disease		
	nephrosis					
	25,399	20,935	96	85	519	422
1						

HIV indicates human immunodeficiency virus.

Note: Deaths within each age group do not sum to all ages combined due to the inclusion of unknown ages. In accordance with the National Center for Health Statistics' cause-of-death ranking, "Symptoms, signs, and abnormal clinical or laboratory findings" and categories that begin with "Other" and "All other" were not ranked.

Source: US Final Mortality Data, 2016, National Center for Health Statistics, Centers for Disease Control and Prevention, 2018.

<sup>*a*</sup> Includes primary and secondary hypertension.

Table 9 presents the number of deaths in 2016 for the 5 leading cancer types by age and sex. Brain and other nervous system tumors are the leading cause of cancer death among men aged younger than 40 years and women aged younger than 20 years, whereas breast cancer leads among women aged 20 to 59 years. Lung cancer leads in cancer deaths among men aged 40 years and older and women aged 60 years and older, causing more deaths in 2016 than breast cancer, prostate cancer, CRC, and leukemia combined. There were approximately 20% more lung cancer deaths in men (80,775) than in women (68,095) in 2016, but this pattern is projected to reverse by 2045 if current smoking trends continue.67 Cervical cancer continues to be the second leading cause of cancer death in women aged 20 to 39 years, causing 9 deaths per week in this age group. This finding underscores the need for increased HPV vaccination uptake in adolescents and guideline-adherent screening in young women. Notably, the percentage of women aged 22 to 30 years who had never been screened for cervical cancer increased between 2000 and 2010.68 In addition, an estimated 14 million screening-aged women (ages 21-65 years) had not been tested in the past 3 years in 2015.69

ALL AGES	<20	20 TO 39	40 TO 59	60 TO 79	≥80
		M	ALE		
ALL SITES	ALL SITES	ALL SITES	ALL SITES	ALL SITES	ALL SITES
314,571	1,100	4,044	49,227	172,243	87,954
Lung &	Brain & ONS	Brain & ONS	Lung &	Lung &	Lung &
bronchus			bronchus	bronchus	bronchus

**Table 9.** Five Leading Causes of Cancer Death by Age and Sex, United States, 2016

80,775	314	538	11,588	49,877	19,095
Prostate	Leukemia	Leukemia	Colorectum	Colorectum	Prostate
30,370	280	526	5,888	14,010	15,535
Colorectum	Bones & joints	Colorectum	Liver <sup>a</sup>	Prostate	Colorectum
27,642	120	487	4,001	13,447	7,250
Pancreas	Soft tissue (including	Non-Hodgkin	Pancreas	Pancreas	Urinary
	heart)	lymphoma			bladder
21,899	87	237	3,747	12,926	5,621
Liver <sup>a</sup>	Non-Hodgkin	Soft tissue (including	Brain & ONS	Liver <sup>a</sup>	Pancreas
	lymphoma	heart)			
17,843	42	232	2,562	10,961	5,099
		FEM	1ALE		
ALL SITES	ALL SITES	ALL SITES	ALL SITES	ALL SITES	ALL SITES
283,467	820	4,653	48,075	140,971	88,944
Lung &	Brain & ONS	Breast	Breast	Lung &	Lung &
bronchus				bronchus	bronchus
68,095	238	1,158	10 405	39 029	19 199
Breast			10,405	33,023	15,155
	Leukemia	Uterine cervix	Lung &	Breast	Breast
	Leukemia	Uterine cervix	Lung & bronchus	Breast	Breast
41,488	Leukemia 219	Uterine cervix 469	Lung & bronchus 9,676	Breast 18,922	Breast 11,002
41,488 Colorectum	Leukemia 219 Bone & joints	Uterine cervix 469 Colorectum	Lung & bronchus 9,676 Colorectum	Breast 18,922 Pancreas	Breast 11,002 Colorectum
41,488 Colorectum 24,644	Leukemia 219 Bone & joints 80	Uterine cervix 469 Colorectum 417	Lung & bronchus 9,676 Colorectum 4,328	Breast 18,922 Pancreas 10,971	Breast 11,002 Colorectum 9,637
41,488 Colorectum 24,644 Pancreas	Leukemia 219 Bone & joints 80 Soft tissue (including heart)	Uterine cervix 469 Colorectum 417 Brain & ONS	Lung & bronchus 9,676 Colorectum 4,328 Ovary	Breast 18,922 Pancreas 10,971 Colorectum	Breast 11,002 Colorectum 9,637 Pancreas
41,488 Colorectum 24,644 Pancreas 20,858	Leukemia 219 Bone & joints 80 Soft tissue (including heart) 80	Uterine cervix 469 Colorectum 417 Brain & ONS 371	Lung & bronchus 9,676 Colorectum 4,328 Ovary 2,777	Breast 18,922 Pancreas 10,971 Colorectum 10,259	Breast 11,002 Colorectum 9,637 Pancreas 7,074
41,488 Colorectum 24,644 Pancreas 20,858 Ovary	Leukemia 219 Bone & joints 80 Soft tissue (including heart) 80 Liver	Uterine cervix 469 Colorectum 417 Brain & ONS 371 Leukemia	Lung & bronchus 9,676 Colorectum 4,328 Ovary 2,777 Pancreas	Breast 18,922 Pancreas 10,971 Colorectum 10,259 Ovary	Breast 11,002 Colorectum 9,637 Pancreas 7,074 Leukemia

ONS indicates other nervous system.

Note: Ranking order excludes category titles that begin with the word "Other."

\* Includes intrahepatic bile duct.

### Cancer Disparities by Socioeconomic Status

Lower socioeconomic status (SES), whether measured at the individual or area level, is associated with numerous health disadvantages and higher mortality across race and ethnicity.70-72 A recent study estimated that approximately one-third (34%) of cancer deaths in Americans aged 25 to 74 years could be averted with the elimination of socioeconomic disparities.72 Notably, socioeconomic deprivation was associated with lower cancer mortality prior to the mid-1980s because of the later development of effective treatment and the historically elevated risk of lung and colorectal cancers among individuals with high SES.73, 74

County-level SES indicators only indirectly reflect individual SES, but are valuable because the county is the smallest geographic unit for which policy is legislated. In addition, countylevel indicators potentially capture some of the complex environmental influences on health. Figure 8 depicts the distribution of county-level poverty by quintile across the United States during 2012-2016, when the overall cancer death rate was approximately 20% higher among residents of the poorest compared with the most affluent counties. Socioeconomic inequalities in cancer mortality widened over the past 3 decades overall, but there is substantial variation by cancer type. Consistent with socioeconomic inequalities for cancer incidence,75 the largest gaps are for the most preventable cancers. For example, cervical cancer mortality among women in poor counties is twice that of women in affluent counties, and lung and liver cancer mortality among men is >40% higher (Table 10). The most striking socioeconomic shift occurred for CRC mortality; rates in men in the poorest counties were approximately 20% lower than those in affluent counties in the early 1970s, but are now 35% higher (Fig. 9). This reversal reflects changes in dietary and smoking patterns that influence CRC risk,73 as well as the slower dissemination of screening and treatment advances among disadvantaged populations.76 A similar crossover occurred earlier for male lung cancer mortality because historically, men of higher SES were much more likely to smoke.73



#### Open in figure viewer PowerPoint

County-Level Poverty (in Percentage) in the United States, 2012 to 2016.

**Table 10.** Change in Cancer Mortality Rates by County-Level Poverty, 1970 to 1974 Versus2012 to 2016

	1970 TO 1974			2012	TO 2016		
			RATE RATIO (95% CI)			RATE RATIO (95% CI)	
	POOR	AFFLUENT	POOR VS AFFLUENT	POOR	AFFLUENT	POOR VS AFFLUENT	
All cancers	-						
Both sexes	199.7	198.8	1.00 (1.00-1.01)	176.7	149.7	1.18 (1.18- 1.19)	
Male	259.0	250.4	1.03 (1.03-1.04)	217.5	177.3	1.23 (1.22- 1.23)	

Female	157.5	164.4	0.96 (0.95-0.97)	147.6	130.2	1.13 (1.13- 1.14)
Brain & ONS						
Both sexes	3.7	4.1	0.90 (0.86-0.93)	4.0	4.6	0.89 (0.86- 0.91)
Male	4.6	4.9	0.93 (0.88-0.97)	4.9	5.6	0.87 (0.84- 0.91)
Female	3.0	3.4	0.87 (0.82-0.92)	3.4	3.7	0.91 (0.87- 0.95)
Breast (female)						
All races	29.0	34.0	0.85 (0.84-0.87)	22.5	19.5	1.16 (1.14- 1.17)
White	28.8	34.4	0.84 (0.82-0.85)	20.9	19.7	1.06 (1.04- 1.09)
Black	30.1	30.8	0.97 (0.90-1.05)	28.8	25.7	1.12 (1.08- 1.16)
Colorectum						
Both sexes	25.5	30.9	0.83 (0.81-0.84)	16.5	12.7	1.30 (1.28- 1.32)
Male	28.6	35.6	0.81 (0.79-0.82)	20.2	14.9	1.35 (1.33- 1.38)
Female	23.3	27.8	0.84 (0.82-0.86)	13.6	10.9	1.25 (1.23- 1.28)
Esophagus						
Both sexes	3.9	3.3	1.19 (1.14-1.24)	4.0	3.9	1.02 (0.99- 1.04)
Male	6.7	5.5	1.20 (1.14-1.26)	7.1	6.9	1.03 (1.00- 1.06)
Female	1.8	1.6	1.15 (1.06-1.25)	1.5	1.5	0.99 (0.93- 1.05)

Leukemia						
Both sexes	8.1	8.3	0.97 (0.94-1.00)	6.4	6.4	1.00 (0.97- 1.02)
Male	10.5	11.1	0.94 (0.91-0.98)	8.6	8.7	0.99 (0.97- 1.02)
Female	6.3	6.4	0.99 (0.95-1.04)	4.8	4.8	1.00 (0.97- 1.04)
Liver & intrahepatic bil duct	e					
Both sexes	3.5	2.8	1.27 (1.21-1.33)	7.7	5.6	1.37 (1.35- 1.40)
Male	4.8	3.8	1.29 (1.21-1.37)	11.5	8.2	1.41 (1.37- 1.44)
Female	2.5	2.0	1.22 (1.13-1.31)	4.5	3.5	1.31 (1.27- 1.36)
Lung & bronchus						
Both sexes	41.2	37.3	1.11 (1.09-1.12)	47.7	37.2	1.28 (1.27- 1.29)
Male	76.3	66.8	1.14 (1.13-1.16)	63.0	44.2	1.42 (1.41- 1.44)
Female	14.2	14.7	0.96 (0.94-0.99)	36.1	32.0	1.13 (1.12- 1.14)
Myeloma						
Both sexes	2.8	2.8	1.00 (0.96-1.05)	3.7	3.1	1.17 (1.14- 1.21)
Male	3.4	3.4	1.00 (0.93-1.07)	4.6	4.0	1.14 (1.09- 1.19)
Female	2.3	2.3	1.01 (0.94-1.08)	3.0	2.5	1.22 (1.17- 1.28)

Non-Hodgkin lympho	ma					
Both sexes	4.9	6.0	0.83 (0.80-0.85)	5.6	5.5	1.02 (1.00- 1.05)
Male	6.2	7.3	0.85 (0.81-0.89)	7.2	7.1	1.02 (0.98- 1.05)
Female	4.0	5.0	0.80 (0.76-0.84)	4.4	4.2	1.03 (0.99- 1.06)
Ovary						
All races	9.0	10.6	0.84 (0.81-0.87)	7.0	7.0	1.00 (0.97- 1.03)
White	9.3	10.8	0.86 (0.83-0.90)	7.3	7.3	1.00 (0.97- 1.04)
Black	7.6	8.5	0.89 (0.77-1.04)	6.4	6.1	1.05 (0.97- 1.14)
Pancreas						
Both sexes	10.8	10.5	1.03 (1.01-1.06)	11.4	10.8	1.06 (1.04- 1.07)
Male	14.3	13.3	1.07 (1.04-1.11)	13.0	12.5	1.04 (1.02- 1.07)
Female	8.3	8.4	0.98 (0.95-1.02)	10.1	9.4	1.07 (1.05- 1.10)
Prostate						
All races	32.6	30.2	1.08 (1.05-1.11)	22.5	17.9	1.26 (1.23- 1.28)
White	28.1	29.9	0.94 (0.91-0.97)	18.2	17.7	1.03 (1.00- 1.05)
Black	51.4	52.8	0.97 (0.90-1.06)	42.9	33.7	1.27 (1.21- 1.34)

Urinary bladder

Both sexes	5.2	5.9	0.87 (0.84-0.91)	4.2	4.3	0.96 (0.94- 0.99)
Male	8.5	10.4	0.82 (0.78-0.86)	7.2	7.6	0.95 (0.92- 0.98)
Female	2.9	3.0	0.97 (0.91-1.04)	2.2	2.1	1.03 (0.98- 1.08)
Uterine corpus						
All races	5.9	5.5	1.08 (1.04-1.13)	5.3	4.6	1.15 (1.11- 1.19)
White	5.2	5.4	0.96 (0.92-1.01)	4.3	4.5	0.96 (0.92- 1.00)
Black	9.0	8.6	1.04 (0.90-1.22)	8.9	8.2	1.08 (1.01- 1.16)
Uterine cervix						
All races	8.9	5.1	1.73 (1.66-1.80)	3.2	1.6	2.00 (1.90- 2.10)
White	6.7	4.9	1.36 (1.30-1.43)	2.9	1.6	1.86 (1.75- 1.98)
Black	16.9	12.4	1.37 (1.22-1.54)	4.3	2.4	1.76 (1.57- 1.99)

95% CI indicates 95% confidence interval; ONS, other nervous system.

"Poor" and "affluent" refer to extreme county-level poverty categories: 21.18% to 53.95% and 1.81% to 10.84%, respectively.

Rates are per 100,000 population and age adjusted to the 2000 US standard population. Rate ratio is the unrounded rate in poor counties divided by the corresponding unrounded rate in affluent counties.



#### Open in figure viewer | PowerPoint

Cancer Mortality Rates by County-Level Poverty, United States, 1970 to 2016. Rates are per 100,000 population and are age adjusted to the 2000 US standard population. County-level poverty was derived from the 2012 to 2016 American Community Survey. "Poor" and "affluent" refer to extreme county-level poverty categories: 21.18% to 53.95% and 1.81% to 10.84%, respectively.

In contemporary times, the prevalence of behaviors that increase cancer incidence and mortality are vastly higher among residents of the poorest counties, including double the prevalence of smoking and obesity compared to residents of the wealthiest counties.70 Poverty is also associated with lower cancer screening prevalence,77 later stage diagnosis,78 and a lower likelihood of optimal treatment. Although lack of health care capacity in economically challenged areas likely contributes to these disparities, some states are home to both the poorest and most affluent counties, suggesting an opportunity for improvement in the distribution of services. Increasing access to care weakens the link between SES and health.79 Numerous states have reduced inequalities through various strategies that removed barriers to prevention, early detection, and treatment.80-82

Socioeconomic inequalities in cancer mortality are small or absent for malignancies that are less amenable to prevention or treatment. For example, mortality for leukemia and non-

Hodgkin lymphoma was equivalent across poverty levels, despite a higher incidence in more affluent counties,75 likely reflecting survival disparities.83-85 Inferior survival among those with low SES is predominantly driven by a later stage of disease at diagnosis and less aggressive treatment.86 Disparities are also minimal or nonexistent for pancreatic and ovarian cancers, for which early detection is lacking and even optimal treatment has a nominal influence on survival. The inequality for prostate cancer mortality was largely confined to black men, even after accounting for Hispanic ethnicity among whites (data not shown). This finding is consistent with previous studies showing a stronger association between SES and prostate cancer mortality among blacks.87, 88 The slight excess mortality for brain/other nervous system tumors and urinary bladder cancer in affluent counties is in agreement with incidence studies and may partly reflect detection bias.75, 89

## Cancer Disparities by Race/Ethnicity

Cancer occurrence and outcomes vary considerably between racial and ethnic groups, largely because of inequalities in wealth that lead to differences in risk factor exposures and barriers to high-quality cancer prevention, early detection, and treatment,90, 91 as discussed in the previous section. Cancer incidence and mortality are generally highest among non-Hispanic blacks (NHBs) and lowest among Asian/Pacific Islanders (Table 11). The overall cancer incidence rate in NHB men during 2011 through 2015 was 84% higher than that in Asian/Pacific Islander men and 9% higher than that in NHW men. Notably, NHB women had 7% lower cancer incidence than NHW women (because of lower rates of breast and lung cancer), but 13% higher cancer mortality. In men and women combined, the black-white disparity in overall cancer mortality has declined from a peak of 33% in 1993 (279.0 vs 210.5 per 100,000 population) to 14% in 2016 (183.6 vs 160.7 per 100,000 population). This progress is largely due to the steep drop in smoking prevalence unique among black teens from the late 1970s through the early 1990s.92

**Table 11.** Incidence and Mortality Rates for Selected Cancers by Race and Ethnicity, UnitedStates, 2011 to 2016

	ALL RACES COMBINED	NON- HISPANIC WHITE	NON- HISPANIC BLACK	ASIAN/PACIFIC ISLANDER	AMERICAN INDIAN/ ALASKA NATIVE <sup>a</sup>	HISPANIC
Incidence, 2011-2015						

All sites	449.8	465.3	463.9	291.7	398.5	346.6
Male	494.8	505.5	549.1	298.9	418.4	377.6
Female	419.3	438.4	407.0	290.3	386.9	329.9
Breast	124.7	130.1	126.5	92.9	100.9	93.0
(female)						
Colon &	39.3	39.0	46.6	30.7	44.4	34.4
rectum						
Male	45.2	44.6	55.2	36.1	49.8	41.7
Female	34.3	34.2	40.7	26.4	40.1	28.8
Kidney &	16.4	16.6	18.4	7.8	23.2	16.2
renal pelvis						
Male	22.2	22.5	25.4	11.1	29.9	21.1
Female	11.4	11.4	13.1	5.1	17.4	12.2
Liver &	8.1	6.7	10.7	13.0	14.8	13.3
intrahepatic						
bile duct						
Male	12.5	10.3	17.6	19.9	20.9	19.7
Female	4.3	3.6	5.2	7.4	9.5	7.8
Lung &	60.5	64.7	63.8	34.9	61.5	30.7
bronchus						
Male	71.3	74.3	85.4	44.5	69.3	39.2
Female	52.3	57.4	49.2	27.8	55.7	24.6
Prostate	109.2	101.7	179.2	56.0	73.1	91.6
Stomach	6.6	5.4	10.3	10.5	8.4	9.7
Male	9.1	7.8	14.1	13.7	11.2	12.5
Female	4.6	3.5	7.7	8.0	6.1	7.7
Uterine	7.6	7.1	9.2	6.0	9.2	9.6
cervix						

Mortality, 2012-2016						
All sites	161.0	165.4	190.6	100.4	148.8	113.6
Male	193.1	197.3	239.8	119.1	178.8	138.2
Female	137.7	141.8	160.4	87.0	126.8	96.4
Breast (female)	20.6	20.6	28.9	11.3	14.5	14.3
Colon & rectum	14.2	14.0	19.4	9.9	15.9	11.2
Male	16.9	16.6	24.5	11.7	19.5	14.4
Female	11.9	11.9	16.0	8.4	13.1	8.8
Kidney & renal pelvis	3.8	3.9	3.7	1.8	5.8	3.5
Male	5.5	5.7	5.6	2.7	8.2	5.0
Female	2.3	2.4	2.3	1.1	3.8	2.3
Liver & intrahepatic bile duct	6.5	5.7	8.6	9.4	10.8	9.3
Male	9.6	8.3	13.6	13.9	14.6	13.3
Female	3.9	3.4	4.8	5.8	7.5	6.0
Lung & bronchus	41.9	45.0	45.6	22.8	35.4	18.3
Male	51.6	54.1	63.9	30.3	42.7	25.3
Female	34.4	37.9	33.3	17.4	29.9	13.1
Prostate	19.2	18.1	39.8	8.6	19.1	15.9
Stomach	3.1	2.4	5.7	5.3	5.2	5.1
Male	4.2	3.3	8.4	6.8	7.0	6.5

Female	2.3	1.7	3.9	4.2	3.7	4.0
Uterine	2.3	2.1	3.6	1.7	2.8	2.6
cervix						

Rates are per 100,000 population and age adjusted to the 2000 US standard population. Nonwhite and nonblack race categories are not mutually exclusive of Hispanic origin.

\* Data based on Indian Health Service Contract Health Service Delivery Areas (CHSDA) counties.

## Geographic Variation in Cancer Occurrence

Tables 12 and 13 show cancer incidence and mortality rates for selected cancers by state. State variation in cancer incidence results from differences in medical detection practices and the prevalence of risk factors, such as smoking, obesity, and other health behaviors. For example, up-to-date HPV vaccination coverage among adolescent (ages 13-17 years) boys and girls ranged widely in 2017, from just 29% in Mississippi to 78% in Rhode Island and the District of Columbia.93 This variation may contribute to future differential patterns in HPVassociated cancers across states.94, 95 Geographic health disparities, which have increased over time,96, 97 often reflect the national distribution of poverty.98 This trend may be exacerbated by widening inequalities in access to health care because of state/territory differences in Medicaid expansion and other initiatives to improve insurance coverage.99, 100

STATE	ALL SITES BREAST COLOR		ECTUM LUNG & BRONCHUS		NON-HODGKIN LYMPHOMA		P			
	MALE	FEMALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	Ν
Alabama	518.5	392.8	120.9	51.5	37.1	89.0	51.6	19.7	13.6	
Alaska	420.2	401.2	124.1	45.7	38.6	65.3	50.1	20.9	13.5	
Arizona	403.6	368.6	112.9	38.6	29.1	54.7	45.0	18.3	13.3	
Arkansas	520.6	401.2	114.7	50.8	37.5	98.7	61.6	20.9	14.6	
California	438.2	382.2	121.6	41.5	31.8	49.2	39.0	22.6	15.2	

 Table 12. Incidence Rates for Selected Cancers by State, United States, 2011 to 2015

Colorado	424.4	380.7	123.5	37.8	30.3	46.9	40.7	20.9	14.2
Connecticut	507.6	448.5	140.2	42.9	33.4	67.9	56.2	26.1	17.3
Delaware	552.2	451.8	133.8	42.7	32.8	82.7	62.8	24.8	17.5
Dist. of Columbia <sup>23</sup>	527.8	444.3	144.6	50.1	38.7	65.4	49.5	22.6	12.9
Florida	462.2	389.9	116.0	42.3	32.1	69.3	51.9	20.9	14.5
Georgia	519.5	409.8	125.2	49.3	35.9	82.9	51.7	22.3	14.7
Hawaii	429.2	399.5	136.1	49.8	35.7	56.8	37.6	21.3	14.0
Idaho	463.0	408.6	122.2	39.6	33.2	56.2	46.5	22.4	15.7
Illinois	508.1	435.7	131.7	51.6	37.6	77.8	57.5	23.6	16.3
Indiana	485.4	423.1	121.7	48.3	38.3	88.1	61.4	22.6	16.0
lowa	513.0	433.3	123.4	51.2	39.3	77.1	53.4	26.5	17.8
Kansas <sup>a</sup>	_	—	_	_	_	_	_	_	_
Kentucky	570.2	468.8	125.0	58.0	42.4	112.8	79.0	24.5	16.5
Louisiana	557.2	415.6	124.1	54.9	40.0	87.6	54.4	23.9	16.6
Maine	496.6	448.4	125.7	41.5	33.9	82.5	64.8	23.2	17.7
Maryland	488.4	418.6	131.7	42.0	33.2	65.2	51.8	20.4	14.7
Massachusetts	485.3	445.1	137.6	41.9	33.1	69.3	60.2	23.4	16.3
Michigan	492.8	419.7	123.4	42.8	33.5	75.2	58.5	24.1	16.6
Minnesota*	507.5	438.7	131.5	43.0	34.1	61.6	50.5	26.9	17.9
Mississippi	543.4	401.6	116.0	57.5	41.1	99.8	56.3	20.3	14.3
Missouri	489.7	424.0	128.2	48.8	35.9	87.9	63.9	22.7	15.3
Montana	467.4	415.3	123.2	43.8	33.0	58.6	53.7	21.8	16.4
Nebraska	493.3	415.4	124.1	49.5	37.4	70.6	50.1	24.7	16.8
Nevada <sup>a</sup>	412.2	377.7	109.4	42.5	32.7	59.0	53.8	17.2	12.6
New	511.4	459.2	143.9	42.5	33.9	70.6	62.9	24.8	17.5

Hampshire									
New Jersey	525.2	447.6	133.4	47.9	37.0	64.3	52.6	26.0	18.2
New Mexico <sup>a</sup>	394.1	364.3	112.4	38.2	28.9	46.1	35.6	17.2	13.6
New York	528.1	445.5	131.3	46.0	35.0	69.1	54.1	26.5	17.8
North Carolina	514.6	418.4	131.0	43.3	32.9	86.3	56.5	21.3	14.3
North Dakota	492.8	412.6	123.7	53.0	38.9	68.4	50.7	21.7	17.0
Ohio	497.9	429.5	126.2	48.3	36.4	82.7	59.4	23.1	15.6
Oklahoma	489.8	409.8	118.4	48.1	36.9	85.7	58.7	22.0	15.1
Oregon	453.8	412.4	124.9	39.8	30.4	61.3	52.4	21.8	15.6
Pennsylvania	524.3	455.2	131.0	49.5	37.0	76.5	56.3	25.9	17.9
Rhode Island	505.5	458.1	135.3	40.4	32.5	78.2	64.2	27.0	18.3
South Carolina	512.3	407.5	128.3	44.6	33.7	84.4	53.5	20.2	13.9
South Dakota	484.6	422.2	134.3	48.9	36.8	67.4	51.7	23.6	15.4
Tennessee	514.8	415.2	122.2	46.3	35.6	94.3	61.7	21.6	14.5
Texas	445.9	370.5	111.7	45.7	31.8	65.5	43.5	21.3	14.6
Utah	439.1	371.4	115.1	34.2	27.6	32.4	23.7	22.6	14.9
Vermont	472.4	434.8	130.4	38.7	33.5	69.9	58.3	26.2	18.4
Virginia	444.4	395.6	127.9	40.3	32.3	69.8	50.6	20.4	14.2
Washington	476.5	425.7	135.3	40.0	32.0	62.8	52.1	24.9	16.3
West Virginia	511.0	442.5	116.3	53.2	41.6	98.4	66.2	22.0	15.9
Wisconsin	497.0	430.7	129.7	42.6	33.1	68.0	54.1	25.5	17.2
Wyoming	428.1	375.1	112.6	39.2	27.9	46.6	43.3	19.8	13.9
Puerto Rico <sup>a</sup>	404.9	319.3	93.2	52.5	35.1	24.7	12.3	17.0	12.8
United States	494.8	419.3	124.7	45.2	34.3	71.3	52.3	22.8	15.6

Rates are per 100,000 population and age adjusted to the 2000 US standard population.

— Data unavailable.

\* Data for these states are not included in the US combined rates because either the registry did not consent or high-quality incidence data were not available for all years during 2011 through 2015 according to the North American Association of Central Cancer Registries (NAACCR).

† Rates are based on cases diagnosed during 2011 through 2014.

<sup>‡</sup> Data for Puerto Rico are not included in the US combined rates for comparability to previously published US rates.

#### Table 13. Mortality Rates for Selected Cancers by State, United States, 2012 to 2016

STATE	ALL	SITES	BREAST	COLORECTUM		LUNG & BRONCHUS		NON-HODGKIN LYMPHOMA		
	MALE	FEMALE	FEMALE	MALE	FEMALE	MALE	FEMALE	MALE	FEMALE	ſ
Alabama	226.0	144.9	21.8	19.8	13.0	70.7	37.7	6.9	4.4	
Alaska	189.5	145.9	19.6	17.2	14.1	50.7	37.9	6.4	4.1	
Arizona	167.8	122.8	19.3	15.1	10.6	39.7	29.2	6.3	4.0	
Arkansas	229.4	151.5	21.6	20.5	13.7	75.5	43.0	7.2	4.3	
California	171.0	126.5	19.8	15.1	11.1	36.4	26.4	6.8	4.2	
Colorado	162.6	120.8	19.0	14.0	10.5	32.4	26.7	6.4	3.7	
Connecticut	175.3	128.4	18.1	13.5	10.0	42.0	32.0	7.1	4.1	
Delaware	202.1	145.9	21.4	16.6	10.7	57.1	39.8	8.3	4.7	
Dist. of Columbia	200.2	155.6	28.3	18.4	13.5	44.3	30.7	6.3	3.3	
Florida	182.0	128.5	19.4	15.7	11.0	49.7	33.2	6.8	4.1	
Georgia	206.6	137.0	21.9	19.1	12.1	59.8	33.3	7.0	4.1	
Hawaii	162.3	113.0	16.2	15.7	10.6	39.6	23.9	6.3	3.5	
Idaho	180.1	132.7	20.4	15.4	10.9	40.1	30.1	7.7	5.1	

Illinois	203.0	146.7	21.9	18.7	12.8	55.2	37.6	7.4	4.4
Indiana	217.8	150.2	21.1	18.2	13.1	66.7	41.6	8.4	4.9
Iowa	200.9	139.7	19.1	17.4	13.1	55.7	35.9	8.4	4.8
Kansas	194.4	141.7	20.3	17.5	12.4	53.3	37.3	7.1	4.9
Kentucky	243.7	165.0	21.6	20.2	13.9	84.5	52.2	8.8	4.6
Louisiana	227.6	151.2	23.2	21.0	14.2	67.6	39.3	8.2	4.5
Maine	207.8	148.7	18.4	15.0	11.7	61.6	41.8	7.5	5.0
Maryland	190.7	140.0	22.2	16.9	11.9	48.6	34.3	6.8	4.1
Massachusetts	187.2	135.4	18.0	14.4	10.9	47.6	35.7	6.6	4.2
Michigan	202.4	147.8	21.3	16.8	12.1	56.6	39.8	8.4	4.9
Minnesota	181.2	132.8	18.1	14.4	11.2	44.0	33.3	7.9	4.7
Mississippi	245.5	155.8	23.4	23.1	15.3	78.3	39.9	7.1	4.0
Missouri	210.8	150.2	21.7	18.2	12.7	65.1	43.2	7.0	4.2
Montana	176.5	135.4	20.0	16.2	11.1	41.5	36.1	7.0	4.3
Nebraska	190.1	136.9	20.3	17.6	13.1	50.4	34.3	7.4	4.3
Nevada	184.3	142.9	21.9	19.4	14.0	47.6	39.6	6.5	3.8
New Hampshire	192.0	141.1	19.5	13.9	11.9	50.3	39.9	7.1	4.5
New Jersey	181.4	136.9	21.8	17.5	12.2	43.6	32.0	7.3	4.2
New Mexico	170.4	122.6	18.8	16.5	10.9	35.1	25.7	5.8	4.0
New York	180.5	133.8	19.9	15.9	11.5	45.6	31.7	7.1	4.2
North Carolina	206.0	138.9	20.9	16.7	11.5	62.4	36.5	7.1	4.2
North Dakota	178.8	128.0	17.5	16.2	11.9	47.3	31.2	6.8	4.7
Ohio	212.9	151.9	22.5	18.9	13.2	62.7	41.1	8.0	4.8
Oklahoma	221.5	154.6	22.6	20.9	14.2	67.0	43.3	8.0	4.9

Oregon	189.4	140.9	20.4	15.6	11.4	46.1	35.9	7.9	4.6
Pennsylvania	203.6	145.0	21.6	18.2	13.0	55.2	35.6	7.8	4.7
Rhode Island	201.0	140.4	18.2	15.9	11.3	56.4	40.4	6.5	4.5
South Carolina	213.9	141.3	21.8	17.7	12.2	61.9	35.5	6.8	4.3
South Dakota	192.8	132.7	19.2	19.9	13.2	51.7	33.5	6.9	4.0
Tennessee	227.7	151.5	22.1	19.1	13.2	73.1	42.6	8.2	4.8
Texas	187.0	129.1	20.0	17.8	11.4	47.5	29.4	7.0	4.3
Utah	148.5	109.5	20.1	13.1	9.6	23.4	15.6	6.7	4.3
Vermont	194.0	141.6	18.1	16.2	12.6	49.8	38.1	7.9	4.6
Virginia	194.0	137.4	21.4	16.8	11.5	53.0	34.0	6.9	4.3
Washington	183.6	135.9	19.6	14.5	10.6	44.9	34.1	7.9	4.5
West Virginia	227.1	161.7	21.9	20.9	16.0	72.6	45.1	7.8	4.9
Wisconsin	193.9	139.1	19.5	15.5	11.5	49.6	34.8	7.8	4.4
Wyoming	166.2	128.2	18.1	15.5	10.3	37.3	31.1	7.0	4.4
Puerto Rico <sup>a</sup>	152.7	94.6	17.9	19.7	12.2	19.8	8.9	4.7	2.6
United States	193.1	137.7	20.6	16.9	11.9	51.6	34.4	7.3	4.4

Rates are per 100,000 population and age adjusted to the 2000 US standard population.

<sup>*a*</sup> Rates for Puerto Rico are for 2011 through 2015 and are not included in the overall US combined rates.

The largest geographic variation in cancer occurrence by far is for lung cancer, reflecting the large historical and continuing differences in smoking prevalence between states.101 For example, lung cancer incidence rates during 2011 through 2015 in Kentucky (113 per 100,000 population in men and 79 per 100,000 population in women), where smoking prevalence continues to be highest, were approximately 3.5 times higher than those in Utah (32 per 100,000 population in men and 24 per 100,000 population in women), where smoking prevalence is lowest. In 2016, 1 in 4 residents of Kentucky and West Virginia were

current smokers compared with 1 in 10 in Utah, Puerto Rico, and California.102

### Cancer in Children and Adolescents

Cancer is the second most common cause of death among children aged 1 to 14 years in the United States, surpassed only by accidents. In 2019, an estimated 11,060 children (birth to 14 years) will be diagnosed with cancer and 1,190 will die from the disease. Benign and borderline malignant brain tumors are not included in the 2019 case estimates because the calculation method requires historical data and these tumors were not required to be reported to cancer registries until 2004.

Leukemia is the most common childhood cancer, accounting for 28% of cases (including benign and borderline malignant brain tumors). Brain and other nervous system tumors, approximately one-quarter of which are benign/borderline malignant, are second most common (26%) (Table 14). The distribution of cancers that occur in adolescents (aged 15 to 19 years) differs somewhat from that in children. For example, brain and other nervous system tumors (21%), greater than one-half of which (58%) are benign/borderline malignant, and lymphoma (20%) are the most common cancers, whereas leukemia accounts for just 13% of cases. Thyroid carcinoma and melanoma of the skin account for 11% and 4%, respectively, of cancers in adolescents, but only 2% and 1%, respectively, in children.

**Table 14.** Case Distribution (2011 Through 2015) and 5-Year Relative Survival (2008 Through2014)<sup>a</sup> by Age and ICCC Type, Ages Birth to 19 Years, United States

	BIRTH	ГО 14	15 TO 19		
	PERCENTAGE OF CASES	5-YEAR SURVIVAL, %	PERCENTAGE OF CASES	5-YEAR SURVIVAL, %	
All ICCC groups combined		83.4		84.6	
Lymphoid leukemia	22%	90.8	7%	73.8	
Acute myeloid leukemia	4%	66.4	4%	64.2	
Hodgkin lymphoma	3%	97.8	12%	96.1	
Non-Hodgkin lymphoma (including Burkitt lymphoma)	5%	90.2	7%	89.1	
Central nervous system neoplasms	26%	72.9	21%	77.9	

Neuroblastoma & other peripheral	6%	80.2	<1%	54.1 <sup>a</sup>
nervous cell tumors				
Retinoblasoma	2%	95.2	<1%	—
Nephroblastoma & other	5%	92.7	<1%	_
nonepithelial renal tumors				
Hepatic tumors	2%	80.4	<1%	52.4 <sup>a</sup>
Hepatoblastoma	1%	84.6	<1%	—
Osteosarcoma	2%	69.6	3%	65.7
Ewing tumor & related bone	1%	77.7	2%	64.3
sarcomas				
Rhabdomyosarcoma	3%	70.3	1%	46.2
Germ cell & gonadal tumors	3%	91.6	11%	92.6
Thyroid carcinoma	2%	99.7	11%	99.2
Malignant melanoma	1%	94.9	4%	94.0

ICCC indicates International Classification of Childhood Cancer.

Survival rates are adjusted for normal life expectancy and are based on follow-up of patients through 2015.

- Statistic could not be calculated due to fewer than 25 cases diagnosed during 2008 to 2014.

\* Benign and borderline brain tumors were excluded from survival calculations, but were included in the denominator for case distribution.

† The standard error of the survival rate is between 5 and 10 percentage points.

The overall cancer incidence rate in children and adolescents has been increasing slightly (by 0.7% per year) since 1975. In contrast, death rates have declined continuously for many decades, from 6.5 per 100,000 population in 1970 to 2.3 per 100,000 population in 2016, an overall reduction of 65% (65% in children and 61% in adolescents). Much of this progress reflects the dramatic 78% decline in leukemia mortality, from 2.7 per 100,000 children and adolescents in 1970 to 0.6 in 2016. Improved remission rates of 90% to 100% for childhood acute lymphocytic leukemia over the past 4 decades have been achieved primarily through

the optimization of established chemotherapeutic agents as opposed to the development of new therapies.103 The 5-year relative survival rate for all cancers combined improved from 58% during the mid-1970s to 83% during 2008 through 2014 for children and from 68% to 85% for adolescents.10 However, survival varies substantially by cancer type and age at diagnosis (Table 14).

# Limitations

Although the estimated numbers of new cancer cases and deaths expected to occur in 2019 provide a reasonably accurate portrayal of the contemporary cancer burden, they are model-based, 3-year- and 4-year-ahead projections that should be interpreted with caution and not be used to track trends over time. First, the estimates may be affected by changes in methodology as we take advantage of improvements in modeling techniques and cancer surveillance coverage. Second, although the models are robust, they can only account for trends through the most recent data year (currently 2015 for incidence and 2016 for mortality) and cannot anticipate abrupt fluctuations for cancers affected by changes in detection practice (eg, PSA testing and prostate cancer). Third, the model can be oversensitive to sudden or large changes in observed data. The most informative metrics for tracking cancer trends are age-standardized or age-specific cancer death rates from the NCHS and cancer incidence rates from SEER, NPCR, and/or NAACCR.

Errors in reporting race/ethnicity in medical records and on death certificates may result in underestimates of cancer incidence and mortality in nonwhite and nonblack populations, particularly American Indian/Alaska Native populations. It is also important to note that cancer data in the United States are primarily reported for broad, heterogeneous racial and ethnic groups, masking important differences in the cancer burden within these populations. For example, lung cancer incidence is equivalent in Native Hawaiian and NHW men, but approximately 50% lower in Asians/Pacific Islanders overall.66

# Conclusions

The continuous decline in cancer death rates since 1991 has resulted in an overall drop of 27%, translating to approximately 2.6 million fewer cancer deaths. Although the racial gap in cancer mortality is slowly narrowing, socioeconomic inequalities are widening, with residents of the poorest counties experiencing an increasingly disproportionate burden of the most preventable cancers. These counties are low-hanging fruit for locally focused cancer control efforts, including increased access to basic health care and interventions for smoking cessation, healthy living, and cancer screening programs. A broader application of existing cancer control knowledge with an emphasis on disadvantaged groups would

undoubtedly accelerate progress against cancer.

## Disclosures

All authors are employed by the American Cancer Society, which receives grants from private and corporate foundations, including foundations associated with companies in the health sector for research outside of the submitted work. The authors are not funded by or key personnel for any of these grants and their salary is solely funded through American Cancer Society funds.



Copyright © 1999-2018 John Wiley & Sons, Inc. All rights reserved

